


PrEP Visit Checklist for Providers

TIMELINE



INDICATIONS

- Weighs at least 35 kg (~77 lbs)
- Without evidence of acute or established HIV infection
- Adequate renal function (creatinine clearance ≥ 60 mL/min)
- Increased risk of HIV acquisition through sex or injection drug use, including one or more of the following for the patient AND/OR their partner(s):
 - Condomless vaginal or anal sex (receptive or insertive) with a partner who is HIV infected or of unknown infection status in the past 6 months
 - An HIV-positive partner with a detectable or unknown viral load
 - Diagnosis of a bacterial STI (gonorrhea, syphilis, chlamydia) in the past 6 months
 - A history of stimulant (such as methamphetamine) or injection drug use
 - A history of transactional sex activity
 - A history of intimate partner violence

LABS

- HIV Test - 4th Gen
- RPR
- GC/CT (3-site, if indicated)
- Hep A, B, and C
- Metabolic Panel
- Urinalysis
- Pregnancy Test (if indicated)
- Assess for Acute HIV Infection

PSYCHOSOCIAL EVALUATION

- Assess prospective adherence issues
- Provide risk-reduction counseling
- Offer condoms
- Manage side effects
- Mental health screening and referral is needed
- Assess reproductive health plans

OTHER

- Provide same day initial PrEP Prescription when possible
- No refills, only 30 days of medication

INITIAL
VISIT

- HIV Test - 4th Gen
- RPR
- GC/CT (3-site, if indicated)
- Metabolic Panel
- Urinalysis
- Pregnancy Test (if indicated)
- Assess for Acute HIV Infection

- Assess adherence
- Provide risk-reduction counseling
- Offer condoms
- Manage side effects
- Mental health screening and referral is needed
- Assess reproductive health plans

- Provide PrEP prescription with 3 refills ONLY
- Provide any needed vaccinations and contraceptive modalities where necessary
- If notable adherence concerns present, patient should return in 30 days**

30-DAY
VISIT

- HIV Test - 4th Gen
- RPR
- GC/CT (3-site, if indicated)
- Hep C (quarterly to annually per risk)
- Metabolic Panel
- Urinalysis
- Pregnancy Test (if indicated)
- Assess for Acute HIV Infection

- Assess adherence
- Provide risk-reduction counseling
- Offer condoms
- Manage side effects
- Mental health screening and referral is needed
- Assess reproductive health plans

- Provide PrEP prescription with 3 refills ONLY
- Provide any needed vaccinations and contraceptive modalities where necessary
- If notable adherence concerns present, patient should return in 30 days**

QUARTERLY
VISIT